Guest Editorial

When the Skin Speaks: Rash Recognition as a Patient Safety Priority

Kanika Khandelwal*

ABSTRACT

Rashes are among the most common clinical findings encountered in inpatient care, yet they remain one of the most frequently underestimated. Often dismissed as benign, allergic, or "dermatological," rashes are instead a critical diagnostic window into systemic disease. Delayed or missed diagnosis of serious cutaneous manifestations represents an under-recognized patient safety issue, reflecting broader gaps in clinical prioritization, diagnostic reasoning, and hospital systems.

In an era increasingly focused on patient safety, quality of care, and diagnostic error, rash-related delays deserve explicit attention. The skin is not a peripheral organ. It is a visible, dynamic, and information-rich diagnostic surface, and failure to carefully assess its findings can result in preventable harm.

Many life-threatening conditions declare themselves initially through the skin. Stevens–Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS), cutaneous vasculitis, meningococcemia, infective endocarditis, and necrotizing soft tissue infections may all present with early or subtle rashes. In such conditions, time to recognition often determines outcomes.

In busy inpatient settings, however, rashes are frequently deprioritized compared with laboratory abnormalities or radiologic findings. When patients present with fever, hypotension, sepsis, or organ dysfunction, skin findings may be relegated to secondary documentation, often described vaguely or without reassessment. This oversight allows critical diagnostic clues to be missed during early stages of illness.

Delayed diagnosis of rashes is rarely the result of a single error. Rather, it reflects predictable system-level vulnerabilities. Rashes are often described imprecisely, evolution over time is inadequately documented, and significant clinical overlap exists between infectious, inflammatory, and drug-related eruptions. Empiric treatments—particularly antibiotics and corticosteroids—may be initiated without timely diagnostic reconsideration. Dermatology consultation and skin biopsy both high-yield diagnostic tools, are frequently underutilized or sought late in the clinical course, reducing their diagnostic value.

The patient safety consequences of delayed rash recognition are substantial. Patients may experience progression to multi-organ dysfunction, prolonged hospitalization, unnecessary exposure to broad-spectrum antibiotics or immunosuppressive therapy, and increased risk of adverse outcomes. From a systems perspective, such delays increase length of stay, healthcare costs, and medicolegal risk. Importantly, many of these harms are preventable with earlier recognition and intervention.

These challenges are particularly relevant in Indian healthcare settings, including state and district hospitals across Rajasthan. High patient volumes, limited access to subspecialty consultation, and constrained resources amplify the impact of diagnostic delays. In such environments, improving bedside clinical recognition becomes even more critical. Early and accurate assessment of rashes can prevent unnecessary investigations, reduce inappropriate treatments, and optimize utilization of limited resources.

Reframing rash recognition as a patient safety priority represents a low-cost, high-impact opportunity for quality improvement. Structured skin examination, routine documentation of rash morphology and progression, early reassessment of treatment response, and lower thresholds for dermatology consultation or biopsy in high-risk presentations can meaningfully reduce diagnostic delays. Educational initiatives and simple inpatient algorithms that highlight red flags—such as mucosal involvement, necrosis, rapid progression, or systemic symptoms—can further support frontline clinicians.

Delayed diagnosis of rashes is not merely a dermatology issue; it is a cross-disciplinary patient safety concern. Addressing it requires cultural change, institutional support, and renewed emphasis on clinical observation as a core diagnostic skill. When recognized early, the skin often provides vital clues to underlying disease. Listening to those signals carefully can prevent harm, improve outcomes, and strengthen the quality of inpatient care.

*Assistant Professor, Department of Medicine, Mayo Clinic, Rochester, MN 55905, USA

Corresponding Author:

Dr. Kanika Khandelwal, Assistant Professor, Department of Medicine, Mayo Clinic, Rochester, MN 55905, USA

Email ID: khandelwal.kanika@mayo.edu

This is an open-access article distributed under the terms of Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND) (http://creativecommons.org/licenses/by-nc-nd/4.0/)